

REPORT DOCUMENTATION PAG 18 REPORT SECURITY CLASSIFICATION AD-A231 328							
1a REPORT SECURITY CLASSIFICATION (U)		15 RES N	4D-A2	231	320	>	
Za SECURITY CLASSIFICATION AUTHORITY		3 DISTE	. 🛶	ner URI			
N A 2b DECLASSIFICATION/DOV/NGRAUMG-SCHEDU							
N A			ion Unlimite				
4 PERFORMING ORGANIZATION REPORT NUMBE	•	5 MONITORING	ORGANIZATION RE	PORT N	JMBER(S)		
	Ser						
6a NAME OF PERFORMING ORGANIZATION	6b OFFICE SYMBOL (If applicable)	7a NAME OF M	ONITORING ORGAI	NOITASIN			
University of California,	N A	Office of Namel Bereich					
San Diego 6c ADDRESS (City, State, and ZIP Code)	I N A	Office of Naval Research 7b ADDRESS (City, State, and ZIP Code)					
Marine Biology Research Division		Resident Representative					
Scripps Institution of Oceanograph	У	University of California, San Diego (Q-043)					
La Jolla, CA 92093	Tot office system	La Jolla, CA 92093-0001 9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER					
1)RGANIZATION	8b OFFICE SYMBOL (If applicable)	Y PROCUREMEN	I INZIKOMENI IDI	ENTIFICA	LIOM MOMB	ER	
Office of Naval Research	ONR	N00014-88-K-0079					
8c AUDRESS (City State, and ZIP Code)	1141350	<u> </u>	FUNDING NUMBER				
Biological Sciences Division, Code 800 N. Quincy St.	: 1141MB	PROGRAM ELEMENT NO	PROJECT NO	TASK NO		ORK UNIT CCESSION NO.	
Arlington, VA 22217-5000		İ	İ	ł	1	•	
11 TITLE (Include Security Classification)		1	1	L			
Biology of Symbioses Between Marine Invertebrates and Intracellular Bacteria							
12 PERSONAL AUTHOR(S)							
Horst Felbeck							
13a TYPE OF REPORT 13b TIME C			ORT (Year, Month,	Day)	S PAGE CO	UNT	
16 SUPPLEMENTARY NOTATION							
2							
17 COSATI CODES)						
FIELD GROUP SUB GROUP	eria; oligonucleotide probes; 168 rRNA sequences;						
	RuBP carboxylase - rate to the Aller				•		
19 ABSTRACT (Continue on reverse if necessary and identify by block number)							
The object of our research was to characterize symbiotic chemoautotrophic bacteria using							
molecular techniques. We are concentrating on the sequencing of their 16S rRNA to establish phylogenetic relationships between symbionts from different invertebrate hosts. In addition, we							
compare genes for ribulose-1, 5-bisphosphate-carboxylase in a number of symbiotic systems using							
molecular probes of various origin.							
DITRIBUTION STATEMENT A							
Approved to: public release:							
Distribution Unitaritied							
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT 21 ABSTRACT SECURITY CLASSIFICATION WINCLASSIFIED/UNLIMITED SAME AS RPT DTIC USERS U							
ZZZZ NAME OF RESPONSIBLE INDIVIDUAL	U 226 TELEPHONE	(Include Area Code	11226 0	FFICE SYMP	OL		
The Market of Repropriet Individual						· ·	

DD Form 1473, JUN 86

Previous editions are obs"'ate

SECURITY CLASSIFICATION OF THIS PAGE

Distribution List for Annual and Final Reports

- 1. Put a cover page (Form DD 1473) on your report and attach a copy of the distribution list. Mail one copy of the report to each person on the contractor subset list attached on which your name appears. The other subset list is for your information only. Please don't forget to attach this distribution list to your report - otherwise the folks below think they have mistakenly received the copy meant for the Molecular Biology Program and forward it to us.
- 2. Mail two copies to (include a DTIC Form 50 with these two copies too)

Administrator Defense Technical Information Center Building 5, Cameron Station

Alexandria, VA 22314

Mail one copy to each of the following: 3.

- \times Dr. Michael Marron (a) ONR Code 1141 Molecular Biology Program 800 N. Quincy Street Arlington, VA 22217-5000
- (b) Administrative Contracting Officer ONR Resident Representative (address varies - see copy of your grant)
- (c) Director, Applied Research Directorate ONR Code 12 800 N. Quincy Street Arlington, VA 22217-5000
- (d) Director Office of Naval Technology Code 22 800 N. Quincy Street Arlington, VA 22217-5000

(e) Director Chemical and Biological Sci Div Army Research Office P. 0. Box 12211 Research Triangle Park, NC 27709

- (f) Life Sciences Directorate Air Force Office of Scientific Research Bolling Air Force Base Washington, DC 20332
- Director (g) Naval Research Laboratory Technical Information Div, Code 2627 Washington, DC 20375

Andesign For	1	; _{
N.B OP N	À	
	;	1
l the section and the leading		:
	-	- !
E7		
De Care or	_	
1.48		:
L		
n .		
1111		



FINAL REPORT ON CONTRACT N00014-88-K-0079; R&T 4412026

PRINCIPAL INVESTIGATOR: Horst Felbeck

CONTRACTOR: University of California

CONTRACT TITLE: Biology of Symbioses between Marine Invertebrates and Intracellular Bacteria

START DATE: 1 January 1988

TIME PERIOD: 1/1/1988 - 12/31/1990

RESEARCH OBJECTIVE: To study the properties of symbiotic bacteria and to characterize them by 16S rRNA sequencing

PROGRESS:

Our research focused mainly on three subjects:

- a) The characterization and sequencing of the gene for ribulose bisphosphate carboxylase (RubisCO) from symbiotic bacteria of various origins,
- b) To continue methods development for 16S rRNA sequencing from symbionts in frozen and badly preserved specimens, and
- c) To use these new techniques to sequence 16s DNA from a variety of symbionts

a) RubisCO

We have cloned the gene coding for RubisCO from the sulfur oxidixing symbiont of the gastropod Alvinochoncha hessleri. Nucleotide sequence analysis of the cloned fragment revealed that the large (rbcL) and small (rbcS) subunits of the symbiont RubisCO were organized similarly to the RubisCO operons of free-living photo- and chemoautotrophic prokaryotes. A comparison of aligned sequences showed that the symbiont rbcL gene shared the highest degree of sequence identity with the cyanobacterium Anabaena (69%) while the nucleotide sequence of small subunit was 61% identical to that of the green alga Chlamydomonas rheinhardti. A partial sequence analysis, which included a portion of the rbcL sequence from a symbiont of the bivalve Solemya reidi, revealed that the two symbiont sequences were highly identical -- 85 % at the nucleotide level and 93% at the amino acid level, suggesting a recent common origin. RubisCO activity was expressed in Escherichia coli transformed with a plasmid carrying the RubisCO fragment of the gastropod symbiont in the proper orientation for transcription off of the plasmid lac promoter. The level of activity suggests proper assembly of this deep-sea RubisCO into the holoenzyme.

A reprint of our paper published in the Proceedings of the National Academy of Sciences is included.

b) Methods for 16S rRNA sequencing

After many delays and problems we finally succeeded to develop a method to amplify the genes for 16s rRNA from mixed (i.e. host and symbiont) DNA extracted from frozen tissues and to sequence these genes directly from the amplification product.

We first extract the samples in Guanidinium isothiocyanate followed by further purification through CsCl gradient centrifugation. After an additional purification step (dissolving and re-precipitating) one obtains a preparation suitable for amplification.

We have had severe difficulties in the past sequencing directly from PCR amplified DNA. Within the last year of this centract, however, we have overcome these problems and are now routinely using this approach for sequencing. We are able to obtain sequence information usable for phylogenetic comparisons within a few days without prior cloning. The basic protocol is to sequence the cleaned PCR product directly using the Sequenase TM (United States Biochemical) protocol with either the PCR primer or other internal primer as sequencing primer. With these primers we can sequence two thirds of a 16s gene, i.e., approximately a thousand base pairs, which allows a very thorough and detailed comparison of genes for 16s rRNA from different symbionts. Using this technique we are currently able to determine 350 to 400 bases from each primer for double stranded DNA.

c) Sequencing of 16S rRNA from symbiotic bacteria

The host species of the symbionts whose sequences we have obtained are listed in Table 1.

A phylogenetic tree based on most of these sequences (and additional reference species) is also attached.

PUBLICATIONS AND REPORTS

Stein, J.L., Haygood, M., and H. Felbeck: Diversity of ribulose bisphosphate carboxylase genes in sulfur - oxidizing symbioses. In: Endocytobiology IV, P. Nardon, V. Gianinazzi-Pearson, A.M. Grenier, L. Margulis, D.C. Smith (eds.), Institut National de la Recherche Agronomique, Paris, 343 - 348 (1990)

Distel, D.L.: Detection, identification and phylogenetic analysis of endosymbiotic bacteria using ribosomal RNA sequences. In: Endocytobiology IV, P. Nardon, V. Gianinazzi-Pearson, A.M. Grenier, L. Margulis, D.C. Smith (eds.), Institut National de la Recherche Agronomique, Paris, 339 - 342 (1990)

Felbeck, H.: Symbiosis of bacteria with invertebrates in the deep sea. In: Endocytobiology IV, P. Nardon, V. Gianinazzi-Pearson, A.M. Grenier, L. Margulis, D.C. Smith (eds.), Institut National de la Recherche Agronomique, Paris, 327 - 334 (1990)

Felbeck, H., and D. L. Distel: Prokaryotic symbionts in marine invertebrates. In: The Prokaryotes, 2nd edition (eds. A. Balows, H.G. Trüper, M. Dworkin, W. Harder, K. H. Schleifer), Springer Verl., in press

Stein, J.L., Haygood, M., and H. Felbeck: Nucleotide sequence and expression of a deep sea ribulose 1,5 bisphosphate carboxylase gene cloned from a chemoautotrophic bacterial endosymbiont. PNAS 87:8850-8854 (1990)

Distel, D.L., DeLong, E., and J.B. Waterbury: Phylogenetic characterization and in situ localization of the bacterial symbiont of shipworms (Teredinidae: Blvalvia) using 16S rRNA sequence analysis and oligonucleotide probe hybridization. Appl. Env. Microbiol., submitted

Distel, D.L., and A.P. Wood: Phylogenetic characterization and comparison of Thiobacillus thyasiris and the bacteria! endosymbiont in the gill of Thyasira flexuosa (Thyasiridae: Bivalvia) by 16S rRNA sequence analysis. Arch. Microbiol, in prep.

Distel, D.L., and H. Felbeck: Analysis of the phylogenetic origins of autotrophic bacterial symbioses in marine bivalves by 16S rRNA sequence analysis. In prep.

TRAINING ACTIVITIES:

Research assistantships for Ute Hentschel, Tristan Darland, and Connie Woolfe (50%, part of the year) and salary for postdoctoral researcher Dr. Daniel L. Distel.

Women or Minorities - 1 (U.H.) Non-citizens - 1 (german, U. H.)


AWARDS

Graduate student Jeffrey Stein has been awarded a NASA graduate student fellowship.

Table 1. Species with symbionts whose 16S rRNA sequences have been determined by us.

Bivalves

Superfamily	Family	Genus/Species	Location			
Lucinacea	Lucinidae	Lucinoma aequizonata	Santa Barbara Basin, CA			
		Codakia orbicularis	Nassau, Bahamas			
		Anodontia philippiana	Mangrove Bay, Bermuda			
		Divaricella quadrisulcata	Waderick Wells, Exumas			
:	Thyasiridae	Thyasira flexuosa	Brest Harbor, France			
Glossacea	Vesicomyidae	Calyptogena sp C1KN7	Kaiko trench, Japan			
· 	1	Calyptogena kaikoi	Kaiko trench, Japan			
		Calyptogena laubieri	Kaiko trench, Japan			
	Į	Calyptogena magnifica	Rose Garden, East Pacific Rise			
		Calyptogena sp	Oregon Subduction Zone			
		Calyptogena sp	Monterey Submarine Canyon			
		Calyptogena elongata	Santa Barbara Channel, CA			
Mytilacea	Mytilidae	Bathymodiolus thermophilus	Rose Garden, East Pacific Rise			
		Bathymodiolus sp	Louisiana Hydrocarbon Seeps			
		Bathymodiolus sp	Florida Escarpment Seeps			
Pholadaceae	Teredinidae	Lyrodus pedicellatus	San Diego, CA			
		Bankia gouldi	Fort Pierce, FL			
		Dicyathifer manni	Australia			
		Teredora malleolus	Massachusetts			
Other Symbiont-Containing Taxa						
Riftiida	Riftiidae	Riftia pachyptila	Rose Garden, East Pacific Rise			
Nematoda	Stilbonematinae	Catanema sp.	Carrie Bow, Belize			
	<u> </u>					
Gastropoda		Alvinochonca hessleri	Marianas Back Arc basin			



The horizontal axis represents evolutionary distance (nucleotide substitutions per sequence position). Also included as reference species are the enteric bacteria E. coli and P. vulgaris and the obligate sulfur chemoautotroph Thiomicrospira (strain L-12). Phylogenetic relationships among known sulfur and methane oxidizing symbiotic bacteria based on comparison of 16S rRNA sequences.